

VOLUME 49

PART 2

MEMOIRS

OF THE

QUEENSLAND MUSEUM

© Queensland Museum

PO Box 3300, South Brisbane 4101, Australia

Phone 06 7 3840 7555

Fax 06 7 3846 1226

Email qmlib@qm.qld.gov.au

Website www.qm.qld.gov.au

National Library of Australia card number

ISSN 0079-8835

NOTE

Papers published in this volume and in all previous volumes of the *Memoirs of the Queensland Museum* may be reproduced for scientific research, individual study or other educational purposes. Properly acknowledged quotations may be made but queries regarding the republication of any papers should be addressed to the Director. Copies of the journal can be purchased from the Queensland Museum Shop.

A Guide to Authors is displayed at the Queensland Museum web site www.qm.qld.gov.au/organisation/publications/memoirs/guidetoauthors.pdf

A Queensland Government Project

Typeset at the Queensland Museum

OCCURRENCE OF MYCOBIOTA IN EASTERN AUSTRALIAN SEA TURTLE NESTS

A.D. PHILLOTT, C.J. PARMENTER AND C.J. LIMPUS

Phillott, A.D., Parmenter, C.J. & Limpus, C.J. 2004 06 30: Occurrence of mycobiota in eastern Australian sea turtle nests. *Memoirs of the Queensland Museum* 49(2): 701-703. Brisbane. ISSN 0079-8835.

Major loggerhead (*Caretta caretta*), green (*Chelonia mydas*), hawksbill (*Eretmochelys imbricata*) and flatback (*Natator depressus*) sea turtle rookeries in eastern Australia were surveyed for the presence of fungi in turtle nests. While the fungi *Fusarium oxysporum*, *F. solani* and *Pseudallescheria boydii* were not universally present, they were isolated from the exterior of failed eggs of all turtle species and at all rookeries. Predominant fungi species varied spatially and temporally. □ *Sea turtle, egg, nest, fungi.*

A.D. Phillott & C.J. Parmenter, School of Biological and Environmental Sciences, Central Queensland University, Rockhampton 4702; C.J. Limpus, Queensland Environmental Protection Agency, PO Box 155, Albert St, Brisbane 4001, Australia; 25 May 2001.

Globally, fungi have been described from the exterior and/or embryonic tissue of several species of sea turtle eggs (Table 1). *Fusarium oxysporum* Schlecht., *F. solani* (Mart.) Sacc. and *Pseudallescheria boydii* Negroni & Fischer have been identified from the exterior of failed eggs in sea turtle nests at Heron Island (Phillott et al., 2001). There have been no studies to determine the identity of fungi from failed eggs on turtle rookeries along a single coastline.

Heron Island (23°26'S, 151°55'E), Wreck Island (23°21'S, 151°57'E), Peak Island (23°20.5'S, 150°56'E), Milman Island (11°10'S, 143°00'E) and Mon Repos Conservation Park (24°48'S, 152°27'E) are major rookeries for the loggerhead (*Caretta caretta*), green (*Chelonia mydas*), hawksbill (*Eretmochelys imbricata*) and flatback (*Natator depressus*) turtles. These were surveyed in the 1996/97-1998/99 turtle nesting seasons (Table 2). Nests were located after hatchling emergence and excavated by hand to determine hatch success. A single unhatched egg that appeared by visual inspection to be colonised by fungus was taken from some nests (Table 2) and swabbed using MW170 TRANSTUBE®, Amies Clear Media and refrigerated at 3-5°C. Fungal swabs were incubated on half-strength Potato Dextrose Agar with 0.05gL⁻¹ chloramphenicol to inhibit bacterial growth and then subcultured onto a range of media (e.g. Potato Dextrose Agar, Nutrient Agar, Carnation Leaf Agar) as required for identification. Identification of species not included by Booth (1971) and Seifert (1996) was confirmed by Professor D.E. Ellis. Voucher specimens were lodged with the Queensland Department of Primary Industries (Accession

Numbers: *F. oxysporum* BRIP 28368, *F. solani* BRIP 28369, *P. boydii* BRIP 28370).

F. oxysporum, *F. solani* and *P. boydii* were isolated from sea turtle nests at all rookeries investigated (Table 2). The proportion of nests containing each fungal species varied between rookeries and, at Heron I. (the only rookery to be sampled in multiple years), between nesting seasons. Most samples were of single isolates, however 6.54% of all swabs revealed mixed cultures (1 *F. oxysporum* + *F. solani*; 2 *F. oxysporum* + *P. boydii*; 4 *F. solani* + *P. boydii*). While individual rookery/species census data often show absence of a particular fungus, all fungi were recorded from all turtle species.

As all 3 of the mycobiota isolated from Heron I. (Phillott et al., 2001) are common soil saprophytes, their presence in nests at other turtle rookeries in eastern Australia is not surprising. The temporal variation in predominant fungi observed at Heron I. (1996: *F. solani*; 1997: *P. boydii*) suggests that spatial differences in species composition cannot be solely attributed to geographic location, as it may equally have been influenced by environmental factors (e.g. hydric and thermal) and/or temporal variation.

A comparison of rainfall distribution and volume (1996: 237mm; 1997: 183mm), and air temperature (\pm SD 1996: max. 29.8 \pm 1.7°C, min. 23.1 \pm 1.6°C; 1997: max. 30.9 \pm 2.0°C, min. 24.0 \pm 1.9°C; beach temperature data not available), differed at Heron I. over the November to February sampling period between the 2 seasons. This may have contributed to the change in dominant nest mycobiota, as *F. solani* is more prevalent in areas with relatively high rainfall and low temperature (Burgess &

TABLE 1. Records of fungi associated with failed sea turtle eggs.

Turtle	Rookery	Isolate	Source
Green <i>Chelonia mydas</i>	-	<i>Aspergillus</i>	Solomon & Baird, 1980
Kemp's Ridley <i>Lepidochelys kempii</i>	-	<i>Fusarium</i>	Wyneken et al., 1988
Leatherback <i>Dermochelys coriacea</i>	St Croix, U.S. Virgin Islands	<i>Fusarium</i>	Eckert & Eckert, 1990
Loggerhead <i>Caretta caretta</i>	Göksu Delta, Turkey	<i>Mucor</i>	Peters et al., 1994
Olive Ridley <i>Lepidochelys olivacea</i>	Nancite, Costa Rica	<i>Allescheria, Aspergillus, Cunninghamella, Gliocladiopsis, Fusarium, Homodendrum, Saksenaea</i>	Mo et al., 1990
	Naranjo, Nancite, Costa Rica	<i>Fusarium, Monosporium</i>	Acuña-Mesén, 1992

Summerell, 1992). The germination rate and hyphal growth of all 3 fungi at varying hydric and thermal microclimatic conditions (experienced within natural sea turtle nests) is under further investigation.

Loggerhead turtle nesting populations in Queensland are in rapid decline (50-80% over 10-15yrs dependent on study site: Limpus & Reimer, 1994). Loggerhead nesting in the south Pacific Ocean occurs almost entirely in the southern Great Barrier Reef (the most significant being at Wreck I.) and Bundaberg coastline (including Mon Repos Conservation Park: Limpus, 1985). However loggerhead nests demonstrate high mortality (and associated fungal presence) on Wreck and Heron Islands when compared with green turtle nests at the same location, or loggerhead turtle nests on the adjacent mainland (Limpus et al., 1983; Phillott, 2002). Determining the role of fungi as an opportunistic contaminant of failed turtle eggs or infectant of living embryonic tissue is, therefore, of great conservation importance.

ACKNOWLEDGEMENTS

Dr K.M. Harrower and S.A.M. Elsmore, Central Queensland University, confirmed identification of the *Fusaria* spp. and Prof. D.E. Ellis, Adelaide Women's and Children's Hospital, confirmed identification of *Pseudallescheria boydii*. We thank Dr J.D. Miller, D.J. Limpus, I. Bell and other Queensland Parks and Wildlife Service personnel for field support. This research was carried out under Central Queensland University Ethical Clearance Certificate No. 95/1-104 with approval through Queensland Turtle Research, QPWS. Support was provided by the Centre for Land and Water Resource Management (CQU) Small Grants Scheme.

LITERATURE CITED

- ACUÑA-MESÉN, R.A. 1992. *Monosporium apiospermum* Saccardo (Fungi, Deuteromycetes), asociado a los huevos de la tortuga marina *Lepidochelys olivacea* (Eschscholtz 1829) en Costa Rica. *Brenesia* 38: 159-162.
- BOOTH, C. 1971. The genus *Fusarium*. (Commonwealth Mycological Institute: Kew).
- BURGESS, L.W. & SUMMERELL, B.A. 1992. Mycogeography of *Fusarium*: survey of *Fusarium* species in subtropical and semi-arid grassland soils from Queensland, Australia. *Mycological Research* 96: 780-784.
- ECKERT, K.L. & ECKERT, S.A. 1990. Embryo mortality and hatch success in *in situ* and translocated leatherback sea turtle *Dermochelys coriacea* eggs. *Biological Conservation* 53: 37-46.
- LIMPUS, C.J. 1985. A study of the Loggerhead Turtle, *Caretta caretta*, in Eastern Australia. Unpubl. PhD thesis, University of Queensland, St Lucia.
- LIMPUS, C.J. & REIMER, D. 1994. The loggerhead turtle, *Caretta caretta*, in Queensland: a population in decline. Pp. 34-54. In James, R. (ed.) Proceedings of the Australian Marine Turtle Conservation Workshop, November 1990. (Australian Nature Conservation Agency: Canberra).
- LIMPUS, C.J., REED, P. & MILLER, J.D. 1983. Islands and turtles. The influence of choice of nesting beach on sex ratio. Pp. 397-402. In Baker, J.T., Sammarco, P.W. & Stark, K.P. (eds) Proceedings of the Inaugural Great Barrier Reef Conference. (JCU Press: Townsville).
- MO, C.L., SALAS, I. & CABALLERO, M. 1990. Are fungi and bacteria responsible for olive ridley's egg loss? Pp. 249-252. In Richardson, T.H., Richardson, J.I. & Donnelly, M. (comps) Proceedings of the 10th Annual Workshop on Sea Turtle Biology and Conservation. NOAA Technical Memorandum NMFS-SEFC-278.
- PETERS, A., VERHOEVEN, K.J.F. & STRIJBOSCH, H. 1994. Hatching and emergence in the Turkish Mediterranean loggerhead turtle, *Caretta caretta*: a natural cause for egg and hatchling failure. *Herpetologica* 50: 369-373.

TABLE 2. Spatial and temporal variation in fungal presence in sea turtle nests.

Rookery, Season	Turtle	# Nests Excavated	# Nests Swabbed	% Nests		
				<i>F. oxysporum</i>	<i>F. solani</i>	<i>P. boydii</i>
Heron I. 1996/97	Green	86	22	22.7	63.6	13.6
	Loggerhead	2	2	0.0	100.0	0.0
Heron I. 1997/98	Green	108	19	5.0	5.0	100.0
	Loggerhead	5	3	0.0	0.0	100.0
Milman I. 1998/99	Green	7	5	40.0	60.0	0.0
	Hawksbill	32	8	12.5	75.0	12.5
Mon Repos 1998/99	Loggerhead	12	5	20.0	100.0	20.0
Peak I. 1998/99	Flatback	44	16	12.5	62.5	25.0
Wreck I. 1998/99	Green	39	16	18.8	62.5	37.5
	Loggerhead	7	4	50.0	50.0	25.0

- PHILLOTT, A.D. 2002. Fungal colonisation of sea turtle nests in Eastern Australia. Unpubl. PhD thesis, Central Queensland University, Rockhampton, Queensland.
- PHILLOTT, A.D., PARMENTER, C.J. & LIMPUS, C.J. 2001. Mycoflora identified from failed green (*Chelonia mydas*) and loggerhead (*Caretta caretta*) sea turtle eggs at Heron Island, Australia. *Chelonian Conservation and Biology* 4: 170-172.
- SEIFERT, K. 1996. FusKey: Fusarium Interactive Key. <http://www.res.agr.ca/brd/fusarium>.
- SOLOMON, S.E. & BAIRD, T. 1980. The effect of fungal penetration on the eggshell of the green turtle. Pp. 434-435. In Brederoo, P. & de Priester, W. (eds) *Proceedings of the Seventh European Congress on Electron Microscopy*. (7th European Congress on Electron Microscopy Foundation: Leiden).
- WYNEKEN, J., BURKE, T.J., SALMON, M. & PEDERSON, D.K. 1988. Egg failure in natural and relocated sea turtle nests. *Journal of Herpetology* 22: 88-96.